Open Access



C₆₀ fullerene as promising therapeutic agent for correcting and preventing skeletal muscle fatigue

Yurij I. Prylutskyy¹, Inna V. Vereshchaka², Andriy V. Maznychenko^{2*}, Nataliya V. Bulgakova², Olga O. Gonchar³, Olena A. Kyzyma^{1,4}, Uwe Ritter⁵, Peter Scharff⁵, Tomasz Tomiak⁶, Dmytro M. Nozdrenko¹, Iryna V. Mishchenko⁷ and Alexander I. Kostyukov²

Abstract

Background: Bioactive soluble carbon nanostructures, such as the C_{60} fullerene can bond with up to six electrons, thus serving by a powerful scavenger of reactive oxygen species similarly to many natural antioxidants, widely used to decrease the muscle fatigue effects. The aim of the study is to define action of the pristine C_{60} fullerene aqueous colloid solution (C_{60} FAS), on the post-fatigue recovering of *m. triceps surae* in anaesthetized rats.

Results: During fatigue development, we observed decrease in the muscle effort level before C_{60} FAS administration. After the application of C_{60} FAS, a slower effort decrease, followed by the prolonged retention of a certain level, was recorded. An analysis of the metabolic process changes accompanying muscle fatigue showed an increase in the oxidative stress markers H_2O_2 (hydrogen peroxide) and *TBARS* (thiobarbituric acid reactive substances) in relation to the intact muscles. After C_{60} FAS administration, the *TBARS* content and H_2O_2 level were decreased. The endogenous antioxidant system demonstrated a similar effect because the *GSH* (reduced glutathione) in the muscles and the *CAT* (catalase) enzyme activity were increased during fatigue.

Conclusions: C_{60} FAS leads to reduction in the recovery time of the muscle contraction force and to increase in the time of active muscle functioning before appearance of steady fatigue effects. Therefore, it is possible that C_{60} FAS affects the prooxidant-antioxidant muscle tissue homeostasis, subsequently increasing muscle endurance.

Keywords: C₆₀ fullerene, Skeletal muscles fatigue, Electrical stimulation, Oxidative stress markers, Antioxidant system

Background

Skeletal muscle fatigue is the defence mechanism against overload and leads to the development of painful muscle sensitivity [1-3]. Muscle fatigue develops after physical activities of varying intensities and often leads to acute pain, which can then lead to various chronic disease states [4, 5]. Muscle fatigue is a result of the products of incomplete oxygen oxidation, such as reactive oxygen species (ROS), including peroxides, free radicals, and oxygen ions [6]. During the course of lipid peroxidation, unsaturated fatty acids are formed from various

*Correspondence: maznychenko@biph.kiev.ua





© The Author(s) 2017. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/ publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

² Department of Movement Physiology, Bogomoletz Institute

of Physiology, Bogomoletz Str. 4, Kiev 01024, Ukraine

Full list of author information is available at the end of the article

In this context, bioactive soluble carbon nanostructures, such as the pristine C₆₀ fullerenes, may be considered potential antioxidants [13]. C_{60} fullerene easily bonds with up to six electrons, can serve as a powerful scavenger of ROS [13-15], and is superior to the majority of natural antioxidants, including vitamins C and E and carotenoids, in regard to its antioxidant capacity. As a result, it prevents oxidative stress dissemination in thymocytes [16] and shows a protective effect following the ischemia-reperfusion injury of skeletal muscle [17]. Additionally, water-soluble pristine C₆₀ fullerenes can penetrate through the plasma membrane of cells [18, 19]. Therefore, the use of C₆₀ fullerenes may have a powerful antioxidant effect on the contractile apparatus of striated muscle, thereby facilitating its functional recovery after experimentally induced fatigue.

The aim of this study was to investigate the effect of water-soluble pristine C_{60} fullerenes on the recovery dynamics of the contractile properties of rat *m. triceps surae* (*TS*) after the development muscle fatigue under conditions of long-term activation.

Methods

Material preparation and characterization

A highly stable reproducible pristine C_{60} fullerene aqueous colloid solution (C60FAS) at a concentration of 0.15 mg/ml was prepared according to a previous protocol [20, 21]. Briefly, for the preparation of $C_{60}FAS$ we used a saturated solution of pure C60 fullerene (purity >99.99%) in toluene with a C_{60} molecule concentration corresponding to maximum solubility near 2.9 mg/ml, and the same amount of distilled water in an open beaker. The two phases formed were treated in ultrasonic bath. The procedure was continued until the toluene had completely evaporated and the water phase became yellow colored. Filtration of the aqueous solution allowed to separate the product from undissolved C_{60} fullerenes. The pore size of the filter during the filtration of the aqueous solution was smaller than 2 µm (Typ Whatmann 602 h1/2). The purity of prepared $C_{60}FAS$ (i.e., the presence/absence of any residual impurities, for example carbon black, toluene phase) was determined by HPLC and GC/MS analysis. The maximal concentration of C₆₀ fullerenes in water 0.15 mg/ml was obtained by this method.

The state of C_{60} fullerenes in aqueous solution was monitored using atomic force microscopy (AFM). Under AFM analysis, the sample was deposited onto a cleaved mica substrate (V-1 Grade, SPI Supplies) by precipitation from an aqueous solution droplet. Sample visualization was performed in semi-contact (tapping) mode (Fig. 1a, b). AFM measurements were performed after the complete evaporation of the solvent. Small-angle neutron scattering (SANS) measurements (Fig. 1c) were carried out at the YuMO small-angle diffractometer at the IBR-2 pulsed reactor (JINR, Dubna, Russia) in the time-of-flight mode with the two-detector setup [22]. Treatment of the raw data was performed by the SAS program [23].

Procedure and experimental groups

Male Wistar rats, weighing 280–350 g, were used in the study. The use of the animals was approved by the Ethics Committee of the Institute and performed in accordance with the European Communities Council Directive of 24 November 1986 (86/609/EEC).

The animals were divided into 4 groups. In the experiments, the *m. triceps surae* fatigue was induced by electrical stimulation of *n. tibialis*. Saline solution (group 1, n = 6) or C₆₀FAS (F-injection) 0.1–0.15 mg/kg (group 2, n = 6) was administered into the left *TS* after the development of fatigue. Then, fatigue of the right TS was induced. The data obtained from the ipsilateral (left) side were considered to be the control values vs. those obtained from the contralateral side. The dose range of 0.1–0.15 mg/kg $C_{60}FAS$ does not present any acute or subacute toxicity in rats [13]. The rats of group 3 (n = 6; animals with fatigue of both TS without any injections) and group 4 (n = 6; intact animals) were used only for biochemical studies. After the experiment, the TS of all animals in all groups were removed for biochemical analysis.

It is important to note that a dose of 0.1–0.15 mg/kg C_{60} FAS applied in our experiments does not present any acute or subacute toxicity in animals: it was significantly lower than the maximum tolerated dose of C_{60} fullerene, which was found to be 5 g/kg both for oral or intraperitoneal administration to rats [13]. No toxic effects or death have been fixed under the action of C_{60} fullerenes after their oral administration to rats in total dosage of 2 g/kg for 14 days [24]. Finally, it was shown [13] that water-soluble C_{60} fullerenes administered intraperitoneally to rats (0.5 mg/kg) were subjected to clearance from the organism within 2–4 days.

The animals in groups 1 and 2 were anaesthetized with ketamine (100 mg/kg "Pfizer", USA) combined with xylazine (10 mg/kg, "Interchemie", Holland), tracheostomized and artificially ventilated (out of necessity). The left and right *TS* muscles were separated from the surrounding tissue, and their tendons were detached at the distal insertions. The *n. tibialis* was separated from the tissue and cut proximally, and all branches of the nerve, except those innervating the *TS*, were cut. This nerve was mounted on a bipolar platinum wire electrode for electrical stimulation. The hindlimb muscles and nerves were covered with paraffin oil in a pool formed from skin flaps.





The *TS* muscle was connected via the Achilles tendon to the servo-control muscle puller. The muscle tension was measured by semi-conductor strain gauge resistors glued on a stiff steel beam mounted on the moving part of a linear motor.

To induce muscle fatigue, 1-3 (30 min duration) series intermittent high-frequency electrical stimulation was used (Fig. 2a), separated by rest intervals of 10-20 min. Each series consisted of trains of 0.2-ms rectangular pulses at a rate of 40/s at 12.4 s duration and separated by 5 s intervals of rest (Fig. 2b). The stimulus current was set to 1.3-1.4 times the motor threshold. Note, if muscle fatigue developed in less than 30 min, stimulation was interrupted (it was predicted that fatigue development occurred when there was a muscle force decrease of more than 50% of the initial data). After the end of the 12.4-s-stimulation, the muscle was stretched, and the change in length had a bell-shaped form (one period of 4 Hz sinusoidal signal with corresponding phase locking) of 3.5 mm amplitude and 2 s duration (Fig. 2b; bottom row). The muscle reaction to the stretches appeared as a tension increase after continuous stimulation. These stretches were applied before the post-stimulation twitches to remove, or at least diminish, the after-effects remaining from the continuous stimulation [1]. The signals (stimulus pulses, muscle tension and other) were sampled via DAC-ADC device (CED Power 1401).

Biochemical experiment

For biochemical analysis, the excised *m. triceps surae* (soleus and gastrocnemius) were rapidly dissected, free of fat and tendon, divided into several portions and stored in liquid N₂. For reduced glutathione (*GSH*) analysis, tissue samples were transferred into a medium containing 1 N perchloric acid (1:10 w/v) and homogenized with a motor-driven Potter–Elvehjem glass homogenizer. The resultant homogenate was centrifuged at 10,000g for 10 min (4 °C). The *GSH* content was spectrophotometrically measured [25]. For the enzyme activity assays and H_2O_2 and lipid peroxidation assays, the muscle samples were thawed and homogenized in 50 mM



phosphate buffer with 2 mM EDTA (pH 7.4) at 4 °C (1:9 w/v). Homogenates were then centrifuged for 15 min at 15,000g (4 °C), and the post mitochondrial supernatant was stored at -70 °C.

Oxidative damage in the tissue was measured using the thiobarbituric acid reactive substances (*TBARS*) assay. *TBARS* were isolated by boiling tissue homogenates for 15 min at 100 °C with thiobarbituric acid reagent (0.5% 2-thiobarbituric acid/10% trichloroacetic acid/0.63 M/ dm³ hydrochloric acid) and measuring the absorbance at 532 nm. The results are expressed as nM *TBARS*/mg protein, using $\varepsilon = 1.56 \times 10^5$ dm³/M¹/cm¹ [26].

The H_2O_2 concentration in the tissue homogenates was measured using the FOX method, which is based on the peroxide-mediated oxidation of Fe²⁺, followed by the reaction of Fe³⁺ with xylenol orange (o-cresolsulphonephthalein 3',3"-bis[methylimino] diacetic acid, sodium salt). This method is extremely sensitive and is used to measure low levels of water-soluble hydroperoxide present in the aqueous phase. To determine the H_2O_2 concentration, 500 μ l of the incubation medium was added to 500 μ l of assay reagent (500 µM ammonium ferrous sulphate, 50 mM H₂SO₄, 200 µM xylenol orange, and 200 mM sorbitol). The absorbance of the Fe³⁺-xylenol orange complex (A_{560}) was detected after 45 min. Standard curves of H_2O_2 were obtained for each independent experiment by adding variable amounts of H_2O_2 to 500 µl of basal medium mixed with 500 µl of assay reagent. Data were normalized and expressed as $\mu M H_2 O_2$ per mg protein [27].

Catalase activity was measured by the decomposition of hydrogen peroxide, determined by a decrease in the absorbance at 240 nm [28].

GSH was determined using Ellman's reagent. One millilitre of supernatant was treated with 0.5 ml of Ellman's reagent (5.5'-dithio-bis-nitrobenzoic acid in abs. ethanol) and 0.4 M Tris HCl buffer with 2 mM EDTA, pH 8.9. The absorbance was read at 412 nm in a spectrophotometer [25].

The protein concentration was estimated using the method of Bradford with bovine serum albumin as a standard. All chemicals were purchased from Sigma, Fluka and Merck and were of the highest purity.

Data analysis

In the electrophysiological part of the study, each stimulation series (30 min) was divided into three equal portions (Fig. 2a), which were averaged (maximum 33 stimulation in one portion). The average value of the first portion was set to 100%, and the other series were normalized in relation to this (for each hindlimb). The peak amplitudes of the front (P_1) and rear of the front (P_2) (maxima amplitudes at the site, duration of 1 s, Fig. 2b) of the muscle strength of each single series (12.4 s) were identified and the difference between P_1 and P_2 (ΔP) was calculated. This difference determines the dynamic component of the muscle force decrease in a short period of continuous stimulation. Mean values (mean \pm SD) of the TS muscle strength before and after F-injection were compared using a two-way statistical analysis of variance (ANOVA). The factors of variation included two conditions, time and the effects of the C₆₀FAS. A Bonferroni post hoc analysis was used to determine the differences between groups. The level of significance was set at p < 0.001.

Biochemical data are expressed as the means \pm SEM for each group. The differences among experimental groups were detected by one-way ANOVA followed by Bonferroni's multiple comparison test. Values of p < 0.05 were considered significant.

Results

Analysis of AFM and SANS data

Because the C_{60} fullerene particle size directly correlates with their biodistribution and toxicity [29, 30], the AFM and SANS studies were performed.

The AFM images (Fig. 1a, b) clearly demonstrate randomly arranged, individual C_{60} fullerenes (0.7 nm in diameter) and their bulk clusters with a height of 1.5– 200 nm. At the same time, some individual C_{60} fullerene aggregates with a height of >200 nm are also seen in the AFM image (Fig. 1b). The results obtained are consistent with the theoretical calculations and experimental measurements [20, 21, 31, 32] and demonstrate the polydispersity of the C_{60} FAS used in our study.

Experimental SANS curve for $C_{60}FAS$ is shown in Fig. 1c. The scattering curve of $C_{60}FAS$ is well described by the form-factor of polydisperse spherical particles. The mean radius of gyration of the particle cross section, R_{σ} , and pair distance distribution function, P(r), were found by using indirect Fourier transformation (IFT) approach [33]. We can calculate the radius of particles, R, present in the C₆₀FAS according to well-known equation $R_g^2 = 0.6R^2$ assuming of homogeneous and spherical of C_{60} fullerene clusters. This conclusion follows from previous experimental data [20, 21] and the estimates of the average cluster density according to the contrastvariation experiments [31, 32, 34]. The data given by this procedure indicate that C60FAS consists of C60 fullerene sphere-like nanoparticles with an average size of ~56 nm that is in a good agreement with above AFM data.

It is known [35, 36] that the permeability and cytochemical behavior of nanoparticles strongly depend on their size and, correspondingly, mass (number) distribution. In this regard, our previous studies [16, 18, 19, 29] clear demonstrate that the used C_{60} fullerene nanoparticles can effectively penetrate through the plasma membrane of cells by passive diffusion or endocytosis (depending on the size) and do not exhibit cytotoxic effects.

Electrophysiological experiments

Changes in the *TS* force reaction under fatigue conditions due to prolonged high frequency stimulation (30 min, 40/s) of the *n. tibialis* for animal groups 1 (before and after administration of saline solution) and 2 (before the application of $C_{60}FAS$) did not significantly differ. The analysis was performed by determining the force level at the beginning (P_1) and end (P_2) of single tetanic contractions and the difference between these values (ΔP), which determines the dynamic component of the force decrease during a short period of continuous stimulation (Fig. 2b). The muscle was considered tired if the amplitude of the single tetanic contractions decreased by more than 50% relative to the initial level. When muscle fatigue was reached, the stimulation was stopped and followed by a 10–20 min rest period. Therefore, in the case of one animal, as a result of muscle fatigue stimulation of the left TS, a 50% fatigue level was reached in approximately 12 min; during the next 4 min of stimulation, it continued to decrease [Fig. 3a (I_L), b(I_L)]. After 10 min of rest, a single tetanic contraction force was slightly restored, but it did not reach the initial muscle activity level and continued to decrease rapidly [Fig. 3a (II_I), $b(II_I)$]. In this case, there was also a simultaneous decrease in the dynamic component of the force drop ΔP . Note that the dynamic component was the most highly expressed at the beginning of the first experimental series and that the P_1 amplitude was higher relative to the P_2 amplitude [Fig. 3b (II_I)]. After tetanic contractions for 1.5-2 min, difference between amplitudes P_1 and P_2 was reduced to zero, with moderate variations both in one and the opposite direction over the additional fatigue stimulation period. Simultaneously with the decrease in ΔP values, there was a constant decrease in the developed force. In the following stimulation series, after a period of rest, the initial amplitude of the dynamic component was usually decreased [Fig. 3b (II₁, III₁)].

When a predetermined level of muscle fatigue was reached, C₆₀FAS (0.1-0.15 mg/kg) was injected intramuscularly [at 45 min after the beginning of fatigue stimulation; Fig. 3a (I_{I}^{F}, II_{I}^{F}) , $b(I_{I}^{F}, II_{I}^{F})$]. At the same time, the dynamic changes in the muscle strength level in response to stimulation reflected the further development of fatigue, and the single contraction forces were reduced rapidly [Fig. $3b(I_I^F)$]. However, F-injection led to the gradual recovery of the isometric force levels (at 32 min after drug application; Fig. 3a $[(II_I^F), b (II_I^F)]$. The appearance of negative ΔP values (P_2 amplitude increase compared to P_1 amplitude) indicated the beginning of the recovery [Fig. 3b (II^F_I), 10th min]. In this series of stimulations, the level of the muscle contraction force was recovered to that developed during the initial stages of fatigue stimulation.

Power reaction of the right *TS* was significantly different from the left *TS*. Notably, the *TS* of the right limb was not previously fatigued before the F-injection (Fig. 3c, d). At 52 min after drug administration, a certain force muscle decrease was observed. In this case, the P_1 amplitude was higher than the P_2 amplitude, as indicated by the increase in ΔP values [Fig. 3c (I^F_R), d (I^F_R)]. However, at 6 min after the beginning of fatigue stimulation, the force developed by the muscle appeared at a certain stationary level, which was held during the experimental series. The difference between the P_1 and P_2 amplitudes disappeared (value of ΔP decreased to zero), which may indicate a constant force level at the time of loading. It



Fig. 3 Strength of the IpsI- and contralateral *m*. *triceps surae* (1s) contraction induced by electrical fatigue stimulation before and after $C_{60}FAS$ injection into the *left TS*: **a**, **c** time protocol registrations of the *left* and *right TS* contraction, respectively (*triangles* indicate the moment the of the $C_{60}FAS$ injection); **b**, **d** amplitude values of the muscle force (P_1) at the beginning of single tetanic contractions (*squares*) and ΔP (the difference between the force values at the beginning and at the end of muscle contraction; *triangles*). The rapid development of fatigue (**a**) (decrease in the muscle strength of more than 50%) led to shortening of the stimulation time (I–III). Designations for I–IV (**a**) and I–III at (**c**) correspond to recordings on (**b**) and (**d**), respectively. Indices: *L*, *R* left and right *TS*; *F* registration of muscle force after the administration of $C_{60}FAS$ into the left *TS*



stimulation (S1, S2, and S3; Fig. 2) before and after (*white* and *grey bars*, respectively) C_{60} FAS administration into the left *m. triceps surae* (*TS*): **a**, **b** the results of two fatigue tests before and after (*white* and *grey bars*, respectively) C_{60} FAS administration into the left *TS* (*open bars*) before C_{60} FAS administration, and right *TS* (*grey bars*) at 52 min after C_{60} FAS administration into the left *TS*. *Asterisks* significant differences (p < 0.001) between the muscle strength during time intervals S₁ and S₃ in one or more series of the stimulation. *I*, *II* successive series of fatigue stimulations (normalisation performed by S₁ in series I). Indices: *L*, *R* left and right *TS*; *F* registration of muscle force after the administration of C_{60} FAS into the left *TS*. *Triangle marks* the moment of C_{60} FAS injection

is significant that the muscle maintained the developed force level for an additional 1 h [Fig. 3c (II_R^F , III_R^F), d(II_R^F , III_R^F)]. For this muscle, the total time of the decrease of the isometric force contraction by 50% was 120 min after drug administration. For comparison, the control duration of the fatigue occurrence period was 42 min.

In Fig. 4a, b, a comparison of the force level changes developed by the left *TS* before (I_L) and after (I^F_L) F-injection in two different experiments is presented. The statistical analysis showed a significant (p < 0.001) force decrease during the series of fatigue stimulation before C_{60} FAS administration [Fig. 4a (I_L), b(I_L)]. After F-injection, a recovery of the muscle force for the first animal [Fig. 4a (I^F_L)] and its holding for the second animal [Fig. 4b]

 (I_L^r)] was observed. At the end of the experimental series, the recovery of the active muscle force response was significant compared to that at the beginning and nearly reached the control values. The analysis showed a significant effect for the factors: drug administration (D) and time (T) after administration and their interaction. The corresponding results of this analysis were as follows: F(D) = 2904.47, p(D) < 0.001, F(T) = 42.420, p(T) < 0.001, F(DxT) = 1350.58, p(DxT) < 0.001 (first animal) and F(D) = 122.80, p(D) < 0.001, F(T) = 1058.29, p(T) < 0.001, p(DxT) = 1287.35, p(DxT) < 0.001 (second animal). The data obtained in all experiments (Fig. 4c-h) indicate that the decrease in the developed force after $C_{60}FAS$ administration (I_L^F , I_R^F) was almost two times



slower than in controls. The maximum significant reduction of the muscle force developed during the entire period of fatigue stimulation was 44% after drug administration, whereas in the control this was 85%. For all experimental animals, similar dynamics of the force level decrease in the control and its more gradual decrease after F-injection were observed.

Biochemical experiments

During long-term stimulation of the muscle, metabolic processes change and are a main factor of muscle fatigue. As a result of the fatigue test, the accumulation of lipid peroxidation secondary products and changes in the levels of antioxidants in the tissue of the fatigued muscle were determined. The data clearly demonstrate the increased level of peroxidation and oxidative stress marker *TBARS* and H_2O_2 after fatigue stimulation (Fig. 5a, b). This increase was significant in relation to the intact muscle (*'norm'*) and was 23% (p < 0.05) for *TBARS* and 38% (p < 0.05) for H_2O_2 . After C_{60} FAS administration into the left *TS*, the *TBARS* concentration was significantly reduced compared to fatigue as follows: 29% (p < 0.05) for the left *TS* and 12% (p < 0.05) for the right one. The H_2O_2 level decreases in comparison to the *'fatigue'* group (by 6% for the left *TS* and 7% for the right one), although the H_2O_2 level remained higher in relation to the intact group (p < 0.05). In turn, in response to such changes in the working muscle, an activation of endogenous antioxidants occurred. During fatigue stimulation, the amount of muscle *GSH* quantitatively increased more than two-fold (p < 0.05) and the activity of the antiperoxide enzyme *CAT* also increased. After C₆₀FAS administration, the *GSH* and *CAT* activities were significantly decreased compared to the group '*fatigue*' by 41.8 and 15.4% for *GSH* and 53 and 43% for *CAT* (p < 0.05) for the left and right *TS*, respectively (Fig. 5c, d).

Discussion

In this study, we investigated changes in the contraction force of the rat *m. triceps surae* under fatigue development before and after C_{60} FAS administration. We did not use a level of stimulation above 40 Hz, and the rest period between the experimental series was 15–20 min [2]. This experimental approach allows us to analyse the nature of the muscle contraction force parameter changes under fatigue stimulation before C60FAS application (into the left TS) and directly after F-injection. A marked decrease in the muscle effort level before C60FAS administration (control) was observe in the all experiments both I_L and II_{I.} stimulation series (Fig. 4a-h). It was the result of modified stimulation pattern action, which was due to the influence of the central and peripheral mechanisms of the development of skeletal muscle fatigue [2]. After intramuscular injection of the C₆₀FAS partial ipsilateral TS muscle recovery was registered in two rats. However, the main finding was observed after the application of C_{60} FAS. Not significant a slower effort decrease, followed by the prolonged retention of a certain level was recorded contralaterally in all animals. Decrease in the muscle contraction force was developed more slowly after C₆₀FAS administration compared to the control. It indicates a deceleration of the fatigue process, and the strength restraint at the constant level for a long time (120 min) indicates an increase in the muscle endurance during such conditions. The data obtained in this study indicate that after drug injection, the time for the TS force maximal level decrease to 44% was 120 min. At the same time in the control, the force level of this muscle during the same period decreased to 85%. We suppose, it was caused by antioxidant effects $C_{60}FAS$ on the fatiguing muscle. The duration of the muscle recovery and its rest periods are also important factors for maintaining efficiency and the normal physiological state of the muscle during dynamic work execution [12]. The dynamic component of the single tetanic contraction is likely a reflection of the interaction of the efficiency of the initial increase of the fast motor unit contractile properties and processes of the fatigue strength reduction [37]. Thus, recovery of muscle strength after F-injection both for the preliminary tired and at fresh muscles indicate, that water-soluble pristine C₆₀ fullerenes can penetrate through the plasma membrane of cells [18, 19] and render of powerful antioxidant effect on the contractile apparatus of striated

muscle, thereby facilitating its functional recovery after experimentally induced fatigue.

Under a moderate external load on the muscle, metabolism occurs aerobically. In the actively contracted muscle, metabolism significantly increases, resulting in the accumulation of secondary oxidation products in muscle fibres, which leads to fatigue development [38]. These metabolic processes are a source of oxygen free radicals and contribute to the intensification of lipid peroxidation processes [39-41]. The presence of such metabolism products prevents the adequate implementation of muscle work and increases the duration of the recovery period. Strenuous exercise and endurance training cause oxidative stress in skeletal muscle and can therefore alter the prooxidant-antioxidant balance [42, 43]. Despite extensive research over the years, the relationship between free radical generation, antioxidant enzymes and exercise in skeletal muscle remains controversial [44, 45]. These discrepancies may be related to differences in exercise mode, intensity, duration of the training program, and muscle fibre type. Skeletal muscles are highly heterogeneous. Each muscle fibre type has distinct metabolic characteristics and oxidative potential as well as antioxidant defence capacity [41]. In our study, as a result of fatigue stimulation in working muscle, there was a significant increase in the secondary products of lipid peroxidation and H_2O_2 compared to the intact (unstimulated muscle) muscle (Fig. 5). During intense (physical activity) contraction, the flow of oxygen through muscle cells is greatly increased. High levels of oxygen uptake (up to 100-fold) can lead to excessive ROS generation and are implicated in fatigue, muscle soreness, and myofibril disruption [45]. Moreover, another potential mechanism involved in the oxidative stress response to high-intensity exercise is the redistribution of blood flow, such as elevated blood flow in the heart, lung, and red slow-twitch muscle fibres, leading to increased mitochondrial respiration, which results in an increase in the production of ROS. We found that long-term electrical stimulation of the muscle induced a significant increase in TBARS and H_2O_2 content that led to an increase of *CAT* activity and GSH content in both fast- and slow-twitch muscle fibres. In this case, after $C_{60}FAS$ administration, the oxygen metabolite concentration was significantly lower. This confirms the previous data regarding the protective effect of $C_{60}FAS$ on the immune and antioxidant systems of the body in various pathologies [15, 46]. The mechanisms of effects of this drug can positively influence the processes of endurance and recovery of the active muscles, inactivating the products of its metabolism.

Increased amounts of *GSH* in the stimulated muscle (without drug administration and after its application) are evidence of the compensatory activation of the endogenous antioxidant systems on the irritant action of sufficient strength (Fig. 5). Many studies showed that during intense stress, there is a significant decrease of reduced GSH and an increased concentration of its oxidative form in the myocardium and *m. soleus* [47, 48]. Simultaneously, contradictory data were obtained in the experiments studying endurance [47, 49]. It was found that under physical activity, the amount of reduced GSH in the *m. gastrocnemius* and *DVL* increase. It is likely that in *m. soleus*, a muscle with a high content of myoglobin, all metabolic and biochemical processes occur under aerobic conditions, which use a large number of mitochondrial enzymes, and the accumulation of oxidized GSSG does not have time to reduce [50]. At the same time, the above mentioned processes in the *m. gastrocnemius* occur anaerobically, in contrast to the *m. soleus*. This causes a slow oxidation process and increases the amount of reduced GSH [51, 52]. Under fatigue, after $C_{60}FAS$ administration, the GSH content was somewhat reduced compared to the "fatigue" state, indicating a reduction in oxidative stress and a normalization of the pro- and antioxidant balance in rat muscle tissue (Fig. 5).

An increase of H_2O_2 during exertion leads to an increase in CAT enzyme activity that has a protective antioxidant function by catalysing the decomposition of hydrogen peroxide to water and oxygen. These results are confirmed by previously obtained data from acute experiments on rats with DVL stimulation [47, 52]. An increase of the enzyme activity in response to exercise was also shown in humans [53]. Moreover, some studies indicate an absence of any changes in CAT concentration in the muscles during physical activity [44, 54, 55]. In fact, several reports demonstrated decreases in catalase activity in both oxidative and mixed fibre limb muscles [56, 57]. In our study, after C₆₀FAS administration under fatigue development, the CAT activity was significantly reduced compared to pure fatigue and remained at the control level. It is hypothesized that C₆₀FAS influence the content and activity of endogenous antioxidants and prevent the occurrence of fatigue in actively contracting muscle, thereby contributing to maintenance of its normal physiological state.

Free radical processes increasing is the main pathogenic factor during skeletal muscles fatigue development [58]. Under significant physical activity there is highly overproduction of free radicals in muscle tissue that intensifies the processes of lipid peroxidation, cell membranes damage and antioxidant enzymes inactivation [59]. The active oxygen metabolites cause direct inhibition of respiratory chain mitochondrial enzymes and reducing the balance of ATP/ADF [59]. The above processes in the background of the lactate accumulation with subsequent development of acidosis and blockage of membrane Ca^{2+} channels lead to a pronounced energy deficit and a significant functional activity reduction of muscle tissue [60].

It is known that application of different nature exogenous antioxidants leads to a significant reduction of fatigue skeletal muscle during intense physical activity and increases the onset time of muscle fatigue under prolonged intense endurance exercise [10, 61, 62]. These data demonstrate the feasibility of using antioxidants to correct the level of oxidative stress in the muscle tissue under extreme influences on the body and its efficiency increasing. Since pristine C_{60} fullerenes, as previously shown in various models in vitro and in vivo [13, 15, 63], actively bind free radicals and display a powerful antioxidant properties of direct action, we can assume that the application of water-soluble C₆₀ fullerenes led to the prooxidant-antioxidant balance normalization in the muscle tissue of rats and helped improve the dynamic parameters of muscle contraction.

Conclusion

The use of C_{60} FAS, even at a low therapeutic dose (0.1–0.15 mg/kg) leads to a reduction in the recovery time of the muscle contraction force (after its complete exhaustion state) on the one hand, and an increase in the time of the muscle active work (endurance) until fatigue development on the other. This result illustrates the effect of C_{60} FAS, along with other possible mechanisms, on prooxidant-antioxidant homeostasis in the muscle tissue of rats.

Abbreviations

 C_{60} FAS: pristine C_{60} fullerene aqueous colloid solution; H_2O_2 : hydrogen peroxide; *TBARS*: thiobarbituric acid reactive substances; *GSH*: reduced glutathione; *CAT*: catalase; ROS: reactive oxygen species; NAD⁺: nicotinamide adenine dinucleotide; HCI: hydrochloric acid; AFM: atomic force microscopy; SANS: small-angle neutron scattering; FOX: ferrous ion oxidation xylenol orange; H_2SO_4 : sulphuric acid; EDTA: ethylenediaminetetraacetic acid; DAC: digital to analogue converter; ADC: analogue to digital converter; ANOVA: analysis of variance; DVL: deep portion of vastus lateralis muscle; GSSG: glutathione disulfide.

Authors' contributions

IVV, AVM and NVB designed and performed the experiments, and the in vitro assays were performed by OOG. UR, PS and OAK were responsible for $C_{60}FAS$ synthesis and characterization. TT helped with preparation of the manuscript and provided funding support. DMN and IVM helped collect and analyze data. YuIP and AIK provided supervision and guidance throughout this work. The manuscript was written through contributions of all authors. All authors read and approved the final manuscript.

Author details

¹ Department of Biophysics, Taras Shevchenko National University of Kyiv, Volodymyrska Str. 60, Kiev 01601, Ukraine. ² Department of Movement Physiology, Bogomoletz Institute of Physiology, Bogomoletz Str. 4, Kiev 01024, Ukraine. ³ Department of Hypoxic States Investigation, Bogomoletz Institute of Physiology, Bogomoletz Str. 4, Kiev 01024, Ukraine. ⁴ Joint Institute for Nuclear Research, Joliot-Curie Str. 6, Dubna, Moscow Region, Russia. ⁵ Institute of Chemistry and Biotechnology, Technical University of Ilmenau, Weimarer Str. 25, 98693 Ilmenau, Germany.⁶ University of Physical Education and Sport, Kazimierza Górskiego Str.1, 80-336 Gdansk, Poland.⁷ Lesia Ukrainka Eastern European National University, Volya Avenue 13, Lutsk 43025, Ukraine.

Acknowledgements

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Ethics approval and consent to participate

The use of the animals was approved by the Ethics Committee of the Institute and performed in accordance with the European Communities Council Directive of 24 November 1986 (86/609/EEC).

Funding

This work was supported by Grant 0024/RSA2/2013/52 from Rozwoj Sportu Akademickiego, POLAND.

Received: 7 November 2016 Accepted: 30 December 2016 Published online: 13 January 2017

References

- Kostyukov AI, Hellström F, Korchak OE, Radovanovic S, Ljubisavljevic M, Windhorst U, Johansson H. Fatigue effects in the cat gastrocnemius during frequency-modulated efferent stimulation. Neuroscience. 2000;92:789–99.
- Kostyukov AI, Kalezic I, Serenko SG, Ljubisavljevic M, Windhorst U, Johansson H. Spreading of fatigue-related effects from active to inactive parts in the medial gastrocnemius muscle of the cat. Eur J Appl Physiol. 2002;86:295–307.
- Ervilha UF, Farina D, Arendt-Nielsen L, Graven-Nielsen T. Experimental muscle pain changes motor control strategies in dynamic contractions. Exp Brain Res. 2005;164:215–24.
- Kadetoff D, Kosek E. The effects of static muscular contraction on blood pressure, heart rate, pain ratings and pressure pain thresholds in healthy individuals and patients with fibromyalgia. Eur J Pain. 2007;11:39–47.
- Schomburg ED, Steffens H, Pilyavskii Al, Maisky VA, Brück W, Dibaj P, Sears TA. Long lasting activity of nociceptive muscular afferents facilitates bilateral flexion reflex pattern in the feline spinal cord. Neurosci Res. 2015;95:51–8.
- Allen DG, Lamb GD, Westerblad H. Skeletal muscle fatigue: cellular mechanisms. Physiol Rev. 2008;88:287–332.
- Aruoma OI. Free radicals, oxidants and antioxidants: trend towards the year 2000 and beyond. In: Aruoma O, Halliwell B, editors. Molecular biology of free radicals in human disease. London: OICA International; 1998. p. 1–28.
- Martarelli D, Pompei P. Oxidative stress and antioxidant changes during a 24-hours mountain bike endurance exercise in master athletes. J Sports Med Phys Fit. 2009;49:122–7.
- 9. Richter C. Biophysical consequence of lipid peroxidation in membranes. Chem Phys Lipids. 1987;44:175–89.
- Mach J, Midgley AW, Dank S, Grant R, Bentley DJ. The effect of antioxidant supplementation on fatigue during exercise: potential role for NAD + (H). Nutrients. 2010;2:319–29.
- 11. Reid MB, Stokić DS, Koch SM, Khawli FA, Leis AA. N-acetylcysteine inhibits muscle fatigue in humans. J Clin Invest. 1994;94:2468–74.
- 12. Harris RC, Sale C. Beta-alanine supplementation in high-intensity exercise. Med Sport Sci. 2012;59:1–17.
- Gharbi N, Pressac M, Hadchouel M, Szwarc H, Wilson SR, Moussa F. C₆₀ fullerene is a powerful antioxidant in vivo with no acute or subacute toxicity. Nano Lett. 2005;5:2578–85.
- 14. Sun T, Xu Z. Radical scavenging activities of alpha-alanine $\rm C_{60}$ adduct. Bioorg Med Chem Lett. 2006;16:3731–4.
- 15. Prylutska SV, Grynyuk II, Matyshevska OP, Prylutskyy Yul, Ritter U, Scharff P. Anti-oxidant properties of C_{60} fullerenes in vitro. Fuller Nanotub Carbon Nanostruct. 2008;16:698–705.

- Prylutska SV, Grynyuk II, Grebinyk SM, Matyshevska OP, Prylutskyy YI, Ritter U, Siegmund C, Scharff P. Comparative study of biological action of fullerenes C₆₀ and carbon nanotubes in thymus cells. Mat Wiss Werkst. 2009;40:238–41.
- Nozdrenko DM, Prylutskyy Yul, Ritter U, Scharff P. Protective effect of water-soluble pristine C₆₀ fullerene in ischemia-reperfusion injury of skeletal muscle. Int J Phys Pathophys. 2014. doi:10.1615/IntJPhysPathophys. v5.i2.10.
- Prylutska S, Bilyy R, Overchuk M, Bychko A, Andreichenko K, Stoika R, Rybalchenko V, Prylutskyy Y, Tsierkezos NG. Water-soluble pristine fullerenes C₆₀ increase the specific conductivity and capacity of lipid model membrane and form the channels in cellular plasma membrane. J Biomed Nanotechnol. 2012;8:522–7.
- Panchuk RR, Prylutska SV, Chumak VV, Skorokhyd NR, Lehka LV, Evstigneev MP, Prylutskyy Yul, Berger W, Heffeter P, Scharff P, et al. Application of C₆₀ fullerene-doxorubicin complex for tumor cell treatment in vitro and in vivo. J Biomed Nanotechnol. 2015;11:1139–52.
- Prylutskyy YI, Petrenko VI, Ivankov OI, Kyzyma OA, Bulavin LA, Litsis OO, Evstigneev MP, Cherepanov VV, Naumovets AG, Ritter U. On the origin of C60 fullerene solubility in aqueous solution. Langmuir. 2014;30(14):3967–70.
- 21. Ritter U, Prylutskyy Yul, Evstigneev MP, Davidenko NA, Cherepanov VV, Senenko AI, Marchenko OA, Naumovets AG. Structural features of highly stable reproducible C_{60} fullerene aqueous colloid solution probed by various techniques. Fuller Nanotubes Carbon Nanostruct. 2015;23:530–4.
- 22. Kuklin Al, Islamov AKh, Gordeliy VI. Two-detector system for small-angle neutron scattering instrument. Neutron News. 2005;16:16–8.
- Soloviev AG, Solovieva TM, Stadnik AV, Islamov AH, Kuklin AI. The upgrade of package for preliminary treatment of small-angle scattering spectra. JINR Commun. 2003;10:2003–86.
- 24. Mori T, Takada H, Ito S. Preclinical studies on safety of fullerene upon acute oral administration and evaluation for no mutagenesis. Toxicology. 2006;225:48–54.
- Sedlak J, Lindsay RH. Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. Anal Biochem. 1968;25:192–205.
- 26. Buege JA, Aust SD. Microsomal lipid peroxidation. Methods Enzymol. 1978;52:302–10.
- Wolff SP. Ferrous ion oxidation in presence of ferric ion indicator xylenol orange for measurement of hydroperoxides. Methods Enzymol. 1994;233:182–9.
- Catalase Aebi H. In: Bergmeyer HU, editor. Methods of enzymatic analysis. New York: Academic Press; 1983. p. 276–86.
- Prylutska SV, Matyshevska OP, Golub AA, Prylutskyy Yul, Potebnya GP, Ritter U, Scharff P. Study of C₆₀ fullerenes and C₆₀-containing composites cytotoxicity in vitro. Mater Sci Eng. 2007;27:1121–4.
- Cataldo F. Solubility of fullerenes in fatty acids esters: a new way to deliver in vivo fullerenes Theoretical calculations and experimental results. In: Cataldo F, Da Ros T, editors. Medicinal chemistry and pharmacological potential of fullerenes and carbon nanotubes, series: carbon materials: chemistry and physics, vol. 1. Netherlands: Springer; 2008. doi:10.1007/978-1-4020-6845-4.
- Prylutskyy YI, Durov SS, Bulavin LA, Adamenko II, Moroz KO, Geru II, Dihor IN, Scharff P, Eklund PC, Grigorian L. Structure and thermophysical properties of fullerene C60 aqueous solutions. Int J Thermophys. 2001;22(3):943–56.
- Prylutskyy YI, Buchelnikov AS, Voronin DP, Kostjukov VV, Ritter U, Parkinson JA, Evstigneev MP. C 60 fullerene aggregation in aqueous solution. Phys Chem Chem Phys. 2013;15(23):9351–60.
- Glatter O. A new method for the evaluation of small-angle scattering data. J Appl Cryst. 1977;10:415–21. doi:10.1107/S0021889877013879.
- Avdeev MV, Khokhryakov AA, Tropin TV, Andrievsky GV, Klochkov VK, Derevyanchenko LI, Rosta L, Garamus VM, Priezzhev VB, Korobov MV, et al. Structural features of molecular-colloidal solutions of C60 fullerenes in water by small-angle neutron scattering. Langmuir. 2004;20:4363–8.
- Johnston HJ, Hutchison GR, Christensen FM, Aschberger K, Stone V. The biological mechanisms and physicochemical characteristics responsible for driving fullerene toxicity. Toxicol Sci. 2010;114:162–82.
- 36. Aschberger K, Johnston HJ, Stone V, Aitken RJ, Tran CL, Hankin SM, Peters SA, Tran CL, Christensen FM. Review of fullerene toxicity and exposure

appraisal of a human health risk assessment, based on open literature. Regul Toxicol Pharmacol. 2010;58:455–73.

- De Luca CJ, Contessa P. Hierarchical control of motor units in voluntary contractions. J Neurophysiol. 2012;107:178–95.
- Casey DP, Joyner MJ. Local control of skeletal muscle blood flow during exercise: influence of available oxygen. J Appl Physiol. 2011;111:1527–38.
- Barclay J, Hansel M. Free radicals may contribute to oxidative skeletal muscle fatigue. Can J Physiol Pharmacol. 1991;69:279–84.
- Diaz PT, She ZW, Davis WB, Clanton TL. Hydroxylation of salicylate by the in vitro diaphragm: evidence for hydroxyl radical production during fatigue. J Appl Physiol. 1993;75:540–5.
- Ji LL. Antioxidants and oxidative stress in exercise. Proc Soc Exp Biol Med. 1999;222:283–92.
- Davies KJ, Quintanilha AT, Brooks GA, Packer L. Free radical and tissue damage produced by exercise. Biochem Biophys Res Commun. 1982;107:1198–205.
- Powers SK, Criswell D, Lawler J, Ji LL, Martin D, Herb RA, Dudley G. Influence of exercise and fiber type on antioxidant enzyme activity in rat skeletal muscle. Am J Physiol. 1994;266:R375–80.
- 44. Ji LL. Exercise and oxidative stress: role of the cellular antioxidant systems. Exerc Sport Sci Rev. 1995;23:135–66.
- Clanton TL, Zuo L, Klawitter P. Oxidants and skeletal muscle function: physiologic and pathophysiologic implications. Proc Soc Exp Biol Med. 1999;222:253–62.
- Didenko G, Prylutska S, Kichmarenko Y, Potebnya G, Prylutskyy Y, Slobodyanik N, Ritter U, Scharff P. Evaluation of the antitumor immune response to C₆₀ fullerene. Mat Wiss Werkst. 2013;44:124–8.
- Leeuwenburgh C, Hollander J, Leichtweis S, Griffiths M, Gore M, Ji LL. Adaptations of glutathione antioxidant system to endurance training are tissue and muscle fiber specific. Am J Physiol. 1997;272:R363–9.
- Ramires PR, Hollander J, Fiebig R, Ji LL. Effects of training and dietary glutathione on liver and muscle glutathione status in rats. Med Sci Sports Exerc. 1999;31:S52.
- Sen CK, Marin E, Kretzschmar M, Hanninen O. Skeletal muscle and liver glutathione homeostasis in response to training, exercise, and immobilization. J Appl Physiol. 1992;73:1265–72.

- Leichtweis S, Leeuwenburgh C, Fiebig R, Parmelee D, Yu XX, Ji LL. Rigorous swim training deteriorates mitochondrial function in rat heart. Med Sci Sports Exerc. 1994;26:S69.
- Lew H, Pyke S, Quintanilha A. Changes in the glutathione status of plasma, liver, and muscle following exhaustive exercise in rats. FEBS Lett. 1985;185:262–6.
- 52. Ji LL, Fu RG. Responses of glutathione system and antioxidant enzymes to exhaustive exercise and hydroperoxide. J Appl Physiol. 1992;72:549–54.
- Sen CK. Oxidants and antioxidants in exercise. J Appl Physiol. 1995;79:675–86.
- Jenkins RR. Free radical chemistry: relationship to exercise. Sports Med. 1988;5:156–70.
- 55. Meydani M, Evans WJ. Free radicals, exercise, and aging. In: Yu B, editor. Free radicals in aging. Boca Raton: CRC Press; 1993. p. 183–204.
- Laughlin MN, Simpson T, Sexton WL, Brown OR, Smith JK, Korthuis RJ. Skeletal muscle oxidative capacity, antioxidant enzymes, and exercise training. J Appl Physiol. 1990;68:2337–43.
- Leeuwenburgh C, Frebig R, Chandwancey R, Ji LL. Aging and exercise training in skeletal muscle: responses of glutathione and antioxidant enzyme systems. Am J Physiol. 1994;267:R439–43.
- Lee KP, Shin YJ, Cho SC, Lee SM, Bahn YJ, Kim JY, Kwon ES, Jeong DY, Park SC, Rhee SG, et al. Peroxiredoxin 3 has a crucial role in the contractile function of skeletal muscle by regulating mitochondrial homeostasis. Free Radical Biol Med. 2014;77:298–306.
- Clarkson PM, Thompson HS. Antioxidants: what role do they play in physical activity and health? Am J Clin Nutr. 2000;72:637–46.
- Grassi B, Rossiter HB, Zoladz JA. Skeletal muscle fatigue and decreased efficiency: two sides of the same coin? Exerc Sport Sci Rev. 2015;43:75–83.
- Ferreira LF, Reid MB. Muscle-derived ROS and thiol regulation in muscle fatigue. J Appl Physiol. 2008;104:853–60.
- Hong SS, Lee JY, Lee JS, Lee HW, Kim HG, Lee SK, Park BK, Son CG. The traditional drug Gongjin-Dan ameliorates chronic fatigue in a forced-stress mouse exercise model. J Ethnopharmacol. 2015;168:268–78.
- Krustic PJ, Wasserman E, Keizer PN, Morton JR, Preston KF. Radical reactions of C₆₀. Science. 1991;254:1183–5.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services

Submit your manuscript at www.biomedcentral.com/submit

• Maximum visibility for your research

